

# A golden lower bound for Property W sets

Urban Larsson,  
Killam postdoc, Dalhousie University, Halifax, Canada,  
Second Joint International Meeting of the Israel Mathematical Union  
and the American Mathematical Society

June 17, 2014

# Table of contents

Prologue

A property of a set of positive integers

Wythoff Nim extensions

# Prologue, Hegarty:

*"An overriding feature of **combinatorial number theory** is that one is interested in properties of general sets of integers rather than of individual ones with a special arithmetical structure"*

# Prologue, Hegarty:

*"An overriding feature of combinatorial number theory is that one is interested in properties of general sets of integers rather than of individual ones with a special arithmetical structure"*

# Prologue, Erdős, Rusza et. al.:

- ▶ There is an infinite set of positive integers  $X$  such that if,

## Prologue, Erdős, Rusza et. al.:

- ▶ There is an infinite set of positive integers  $X$  such that if,
- ▶ for all  $x, y, z, w \in X$ ,  $y - x = z - w$  implies  $y = z$  and  $x = w$ ,

## Prologue, Erdős, Rusza et. al.:

- ▶ There is an infinite set of positive integers  $X$  such that if,
- ▶ for all  $x, y, z, w \in X$ ,  $y - x = z - w$  implies  $y = z$  and  $x = w$ ,
- ▶ then

$$\limsup |X \cap \{1, 2, \dots, n\}| \sim n^{\sqrt{2}-1-o(1)}$$

## Prologue, Erdős, Rusza et. al.:

- ▶ There is an infinite set of positive integers  $X$  such that if,
- ▶ for all  $x, y, z, w \in X$ ,  $y - x = z - w$  implies  $y = z$  and  $x = w$ ,
- ▶ then

$$\limsup |X \cap \{1, 2, \dots, n\}| \sim n^{\sqrt{2}-1-o(1)}$$

- ▶ It is also known that, for all  $X$  satisfying this property,  
 $|X \cap \{1, 2, \dots, n\}| \leq \sqrt{n} + O(n^{1/4})$



## Prologue, Erdős, Rusza et. al.:

- ▶ There is an infinite set of positive integers  $X$  such that if,
- ▶ for all  $x, y, z, w \in X$ ,  $y - x = z - w$  implies  $y = z$  and  $x = w$ ,
- ▶ then

$$\limsup |X \cap \{1, 2, \dots, n\}| \sim n^{\sqrt{2}-1-o(1)}$$

- ▶ It is also known that, for all  $X$  satisfying this property,  
 $|X \cap \{1, 2, \dots, n\}| \leq \sqrt{n} + O(n^{1/4})$
- ▶ These type of estimates often concern bounds on the upper asymptotic density of sets, given certain avoidance criteria

# The $A$ and $B$ sets

Let  $A$  denote any infinite set of positive integers. Let  $B$  denote its complement intersected with the positive integers. Then  $A$  and  $B$  are complementary sets on the positive integers. That is  $A \cup B = \mathbb{N}$  and  $A \cap B = \emptyset$ .

# The $A$ and $B$ sequences

We identify the set  $A$  with the unique sequence  $A = (a_n)_{n=1}^{\infty}$  of strictly increasing positive integers. We are looking for an ordering of the elements in  $B$  that, together with the given  $A$ -sequence, satisfies a certain Property W.

# Property W for a pair of sequences

- (1) Suppose that there is an ordering of the elements in  $B$  such that  $\delta_n := b_n - a_n > 0$ , for all  $n$

## Property W for a pair of sequences

- (1) Suppose that there is an ordering of the elements in  $B$  such that  $\delta_n := b_n - a_n > 0$ , for all  $n$
- (2) The pair of sequences  $(A, B)$  satisfies Property W, if (1) holds and in addition, for all  $i, j \in \mathbb{N}$ ,  $\delta_i = \delta_j$  implies  $i = j$

# Property W for a set

The set  $A$  satisfies **Property W** if (2) holds; that is, if  $A$ 's complementary set  $B$  is sufficiently distanced from  $A$  in this precise sense.

## A W-impossible case

$\delta_n$	0	1	2	2	3	5	6	...
$b_n$	0	2	5	6	10	13	15	...
$a_n$	0	1	3	4	7	8	9	...
$n$	0	1	2	3	4	5	6	...

Given the first few elements of the set  $A = \{a_i\}_{i>0}$ , there is **no** ordering of the elements in  $B$ , satisfying Property W. (For later use, let  $b_0 = a_0 = 0$ .)

## How dense must a W-possible A-set be?

$\delta_n$	0	1	2	3	4	5	6	...
$b_n$	0	2	5	7	10	13	15	...
$a_n$	0	1	3	4	6	8	9	...
$n$	0	1	2	3	4	5	6	...

Let  $\phi$  denote the golden ratio. Wythoff Nim's upper P-positions are  $(0, 0), (1, 2), \dots, (a_n, b_n), \dots$ , where for all  $n \in \mathbb{N}$ ,  $a_n = \lfloor \phi n \rfloor$  and  $b_n = \lfloor \phi^2 n \rfloor$ . The consecutive differences  $\delta_n = b_n - a_n$  are the natural numbers in strictly increasing order, that is  $\delta_n = n$  for all  $n$ . Hence  $\{a_i\}$  satisfies Property W.



## $(1, 2)$ -GDWN produces interesting sequences

- ▶ 2-player impartial games: Nim is a famous normal play heap game, alternating play. Take any number of tokens from precisely one heap, at most the whole heap, finitely many heaps. A player who cannot move loses.

# $(1, 2)$ -GDWN produces interesting sequences

- ▶ 2-player impartial games: Nim is a famous normal play heap game, alternating play. Take any number of tokens from precisely one heap, at most the whole heap, finitely many heaps. A player who cannot move loses.
- ▶ Wythoff Nim's moves are as in 2 heap Nim, or instead remove the same number from each heap.

## (1,2)-GDWN produces interesting sequences

- ▶ 2-player impartial games: Nim is a famous normal play heap game, alternating play. Take any number of tokens from precisely one heap, at most the whole heap, finitely many heaps. A player who cannot move loses.
- ▶ Wythoff Nim's moves are as in 2 heap Nim, or instead remove the same number from each heap.
- ▶ (1,2)-GDWN's rules are: move as in Wythoff Nim, or instead remove  $t > 0$  tokens from one heap and  $2t$  tokens from the other, only limited by the number of tokens in each heap.

## The initial P-positions of (1, 2)-GDWN

$\delta_n$	0	2	4	1	3	6	8	...
$b_n$	0	3	6	5	10	14	17	...
$a_n$	0	1	2	4	7	8	9	...
$n$	0	1	2	3	4	5	6	...


Note that neither  $(b_n)$  nor  $(\delta_n)$  is increasing. By the rules of game it follows that  $\{a_i > 0\} \cup \{b_i > 0\} = \mathbb{N}$ ,  $\{a_i > 0\} \cap \{b_i > 0\} = \emptyset$ , and property W holds.

## Comparing the entries of lower sequences


$x_n$	0	1	3	4	7	8	9	...
$a_n$	0	1	2	4	7	8	9	...
$A_n$	0	1	3	4	6	8	9	...
$n$	0	1	2	3	4	5	6	...

The  $x_n$  entries represent our W-impossible lower sequence,  $a_n$  GDWN and  $A_n$  Wythoff Nim. Ah, they look so similar! How can we distinguish some interesting behavior?


## Detour: moves and P-positions in the first quadrant



Nim's moves,  $(0, t), (t, 0), t \in \mathbb{N}$




# Nim's moves and its single P-beam







# Wythoff Nim's moves




# Wythoff Nim's moves and its splitted P-beams




# (1, 2)-GDWN's moves




# (1,2)-GDWN's moves and P-beams experimentally




# $(1, 2)$ -GDWN's sequence of $b_i/a_i$




# $(1, 2)$ -GDWN's lower subsequence of $b_i/a_i$



# $(1, 2)$ -GDWN's upper subsequence $b_i/a_i$



# $\Theta_m$ : (1, 2)-GDWN's upper P-beams (2, 0.05)-split





# Wythoff Nim extensions and Property W

- ▶ The result on the previous slide is made possible by bounding the lower asymptotic density of any  $a$ -sequence satisfying Property W.

# Wythoff Nim extensions and Property W

- ▶ The result on the previous slide is made possible by bounding the lower asymptotic density of any  $a$ -sequence satisfying Property W.
- ▶ A game is a **Wythoff Nim extension**, if we can define its set of P-positions as  $\{(a_i, b_i), (b_i, a_i)\}$ , with  $(a_i)$  increasing,  $\{a_i\}$  and  $\{b_i\}$  complementary, and such that  $\{a_i\}$  satisfies Property W.

# Wythoff Nim extensions and Property W

- ▶ The result on the previous slide is made possible by bounding the lower asymptotic density of any  $a$ -sequence satisfying Property W.
- ▶ A game is a **Wythoff Nim extension**, if we can define its set of P-positions as  $\{(a_i, b_i), (b_i, a_i)\}$ , with  $(a_i)$  increasing,  $\{a_i\}$  and  $\{b_i\}$  complementary, and such that  $\{a_i\}$  satisfies Property W.
- ▶ **Observation:** The game  $(1, 2)$ -GDWN is a Wythoff Nim extension.

## Why a split? Explanation of Detour

### Lemma

Consider  $(1, 2)$ -GDWN. Suppose, for  $n \in \mathbb{N}$ ,

$$\frac{\#\{i > 0 \mid a_i < n\}}{n} \geq \phi^{-1} - o(1).$$

Then the upper  $P$ -positions split.

## Take a larger view

### Theorem (Property W)

Suppose that  $\{a_i\}$  satisfies Property W. Then, for  $n \in \mathbb{N}$ ,

$$\frac{|\{i > 0 \mid a_i < n\}|}{n} \geq \phi^{-1} - o(1) \quad (1)$$

and

$$\frac{|\{i > 0 \mid b_i < n\}|}{n} \leq \phi^{-2} + o(1). \quad (2)$$

In particular the result holds for  $\{(a_i, b_i)\}$  representing the upper  $P$ -positions of any Wythoff Nim extension.

# Proof

- ▶ Define the  $y$ -sequence as the unique permutation of a given  $b$ -sequence, with entries in increasing order. That is  $y_n < y_{n+1}$  for all  $n$  and  $\{y_n\} = \{b_n\}$ .

## Proof

- ▶ Define the  $y$ -sequence as the unique permutation of a given  $b$ -sequence, with entries in increasing order. That is  $y_n < y_{n+1}$  for all  $n$  and  $\{y_n\} = \{b_n\}$ .
- ▶ Define the unique surjective index-function  $j : \mathbb{N} \rightarrow \mathbb{N}$ ,  $j = j(n)$  such that, for all  $n$ ,  $a_j \leq n < a_{j+1}$ . (This is well defined by  $(a_i)$  strictly increasing and  $a_1 = 1$ .)

- ▶ Suppose that (1) does not hold.



- ▶ Suppose that (1) does not hold.
- ▶ Then, by  $(a_i)$  increasing, there is an  $\epsilon' > 0$  such that, for all sufficiently large  $n$ ,  $\frac{j(n)}{a_{j(n)}} < \phi^{-1} - \epsilon'$ .

- ▶ Suppose that (1) does not hold.
- ▶ Then, by  $(a_i)$  increasing, there is an  $\epsilon' > 0$  such that, for all sufficiently large  $n$ ,  $\frac{j(n)}{a_{j(n)}} < \phi^{-1} - \epsilon'$ .
- ▶ We get  $\frac{1}{\phi^{-1} - \epsilon'} < \frac{a_{j(n)}}{j(n)}$ , which implies that there is an  $\epsilon > 0$  such that, for all sufficiently large  $n$ ,  $\phi n + \frac{\epsilon n}{2} < a_n$ .

- ▶ Suppose that (1) does not hold.
- ▶ Then, by  $(a_i)$  increasing, there is an  $\epsilon' > 0$  such that, for all sufficiently large  $n$ ,  $\frac{j(n)}{a_{j(n)}} < \phi^{-1} - \epsilon'$ .
- ▶ We get  $\frac{1}{\phi^{-1} - \epsilon'} < \frac{a_{j(n)}}{j(n)}$ , which implies that there is an  $\epsilon > 0$  such that, for all sufficiently large  $n$ ,  $\phi n + \frac{\epsilon n}{2} < a_n$ .
- ▶ By complementarity this implies, for all sufficiently large  $n$ ,  $\phi^2 n - \gamma(\epsilon) \geq y_n$ , where  $\gamma(\epsilon) > \frac{\epsilon n}{2}$  is a function of  $\epsilon$  only.

Thus

$$\begin{aligned}\delta'_n &:= y_n - a_n \\ &< (\phi^2 - \phi)n - \epsilon n \\ &= (1 - \epsilon)n,\end{aligned}$$

for all sufficiently large  $n$ . Hence,  $(\delta'_n)_{n \leq N}$  must contain at least  $\epsilon N$  (pairwise) repetitions, for all sufficiently large  $N$ .

Thus

$$\begin{aligned}\delta'_n &:= y_n - a_n \\ &< (\phi^2 - \phi)n - \epsilon n \\ &= (1 - \epsilon)n,\end{aligned}$$

for all sufficiently large  $n$ . Hence,  $(\delta'_n)_{n \leq N}$  must contain at least  $\epsilon N$  (pairwise) repetitions, for all sufficiently large  $N$ . **But this does not yet contradict Property W.** We must show that for any  $b$ -sequence, some  $\delta$ -repetition will be forced.

- ▶ Given  $C \in \mathbb{N}$ , define the finite set  $S_b = S_b(C)$  of all indices of  $b$ -entries smaller than  $C$ .

- ▶ Given  $C \in \mathbb{N}$ , define the finite set  $S_b = S_b(C)$  of all indices of  $b$ -entries smaller than  $C$ .
- ▶ Let  $(n_i)$  be the unique increasing sequence of the numbers in  $S_b$ .

- ▶ Given  $C \in \mathbb{N}$ , define the finite set  $S_b = S_b(C)$  of all indices of  $b$ -entries smaller than  $C$ .
- ▶ Let  $(n_i)$  be the unique increasing sequence of the numbers in  $S_b$ .
- ▶ Then,  $n_i \geq i$ , for all  $i$ , and therefore also, by  $(a_i)$  increasing,  $a_{n_i} \geq a_i$ , for all  $i$ .



- ▶ Suppose now that  $N$  is sufficiently large, so that  $(\delta'_n)_{n \leq N}$  contains  $\epsilon N$  repetitions, as defined in the previous paragraph, and study the unique set  $S_b$  of size  $N$ .

- ▶ Suppose now that  $N$  is sufficiently large, so that  $(\delta'_n)_{n \leq N}$  contains  $\epsilon N$  repetitions, as defined in the previous paragraph, and study the unique set  $S_b$  of size  $N$ .
- ▶ It contains the indices of the  $N$  smallest entries in the  $b$ -sequence,  $(n_i)_{i=1}^N$ .


- ▶ Suppose now that  $N$  is sufficiently large, so that  $(\delta'_n)_{n \leq N}$  contains  $\epsilon N$  repetitions, as defined in the previous paragraph, and study the unique set  $S_b$  of size  $N$ .
- ▶ It contains the indices of the  $N$  smallest entries in the  $b$ -sequence,  $(n_i)_{i=1}^N$ .
- ▶ Thus, since  $\sum_{S_b} a_i \geq \sum_{i=1}^N a_i$  and  $\sum_{S_b} b_i = \sum_{i=1}^N y_i$ , we get  $\sum_{S_b} \delta_i \leq \sum_{i=1}^N \delta'_i$ , and so the sequence  $(\delta_i)_{i=1}^N$  must also contain at least  $\epsilon N$  repetitions for all sufficiently large  $N$ .

- ▶ Suppose now that  $N$  is sufficiently large, so that  $(\delta'_n)_{n \leq N}$  contains  $\epsilon N$  repetitions, as defined in the previous paragraph, and study the unique set  $S_b$  of size  $N$ .
- ▶ It contains the indices of the  $N$  smallest entries in the  $b$ -sequence,  $(n_i)_{i=1}^N$ .
- ▶ Thus, since  $\sum_{S_b} a_i \geq \sum_{i=1}^N a_i$  and  $\sum_{S_b} b_i = \sum_{i=1}^N y_i$ , we get  $\sum_{S_b} \delta_i \leq \sum_{i=1}^N \delta'_i$ , and so the sequence  $(\delta_i)_{i=1}^N$  must also contain at least  $\epsilon N$  repetitions for all sufficiently large  $N$ .
- ▶ This contradicts property W, and so (1) must hold, and thus, by complementarity also (2). □


The other lemma is nontrivial, but very specific for  $(1, 2)$ -GDWN (2014 in JIS). In this talk I just wanted to emphasize the more general result for Property W, which I think will find more interesting applications in the future.

The other lemma is nontrivial, but very specific for  $(1, 2)$ -GDWN (2014 in JIS). In this talk I just wanted to emphasize the more general result for Property W, which I think will find more interesting applications in the future. At last, some plots of similar game's P-positions, some of which exhibit log-periodic behavior along split P-beams and others 'just' split, yet others just distort Wythoffs P-beams.. (mostly conjectures):

## $(2, 3)$ -GDWN sequence $b_i/a_i$




# $(2, 3)$ -GDWN lower subsequence $b_i/a_i$







## $(2, 3)$ -GDWN upper subsequence $b_i/a_i$




$(2, 4)$ -GDWN sequence  $b_i/a_i \rightarrow \phi$ ?




$(p, q)$ -GDWN sequence for non-Wythoff pairs:  $B_i/A_i \rightarrow \phi$




# $(3, 5)$ -GDWN sequence $b_i/a_i$




# $(3, 5)$ -GDWN lower subsequence $b_i/a_i$




# $(3, 5)$ -GDWN upper subsequence $b_i/a_i$



# (4, 6)-GDWN sequence $b_i/a_i$




# $(4, 6)$ -GDWN lower subsequence $b_i/a_i$







# (4, 6)-GDWN upper subsequence $b_i/a_i$




# $(4, 7)$ -GDWN sequence $b_i/a_i$




# (4, 7)-GDWN lower subsequence $b_i/a_i$




# $(4, 7)$ -GDWN upper subsequence $b_i/a_i$



# (731, 1183)-GDWN sequence $b_i/a_i$ (a Wythoff pair)



# (731, 1183)-GDWN sequence $b_i/a_i$ (a Wythoff pair)



# P-beams split for $(1, 2)(2, 3)(3, 5)(5, 8)$ -GDWN?

